

## 2008 SOR guidelines for the prevention and treatment of thrombosis associated with central venous catheters in patients with cancer: report from the working group

P. Debourdeau<sup>1\*</sup>, D. Kassab Chahmi<sup>2</sup>, G. Le Gal<sup>3</sup>, I. Kriegel<sup>4</sup>, E. Desruennes<sup>5</sup>, M.-C. Douard<sup>6</sup>, I. Elalamy<sup>7</sup>, G. Meyer<sup>8</sup>, P. Mismetti<sup>9</sup>, M. Pavic<sup>1</sup>, M.-L. Scrobohaci<sup>10</sup>, H. Lévesque<sup>11</sup>, J. M. Renaudin<sup>12</sup> & D. Farge<sup>13</sup> on behalf of the working group of the SOR

<sup>1</sup>Department of Oncology and Internal Medicine, Desgenettes Hospital, Lyons; <sup>2</sup>SOR, National Cancer Institute, Boulogne-Billancourt; <sup>3</sup>Department of Internal Medicine, La Cavale-Blanche Hospital, Brest; <sup>4</sup>Department of Anesthesiology, Curie Institute, Paris; <sup>5</sup>Department of Anesthesiology, Gustave Roussy Institute, Villejuif; <sup>6</sup>Department of Anesthesiology, Saint Louis Hospital, Paris; <sup>7</sup>Hemostasis Laboratory, Tenon Hospital, Paris; <sup>8</sup>Department of Pneumology, Georges Pompidou Hospital, Paris; <sup>9</sup>Department of Vascular Pathology, Saint-Etienne Hospital, Saint-Étienne; <sup>10</sup>Hemostasis Laboratory, Saint-Louis Hospital, Paris; <sup>11</sup>Department of Vascular Pathology, Bois Guillaume Hospital, Rouen; <sup>12</sup>Department of Vascular Pathology, Georges Pompidou Hospital, Paris and <sup>13</sup>Department of Vascular Pathology, Saint-Louis Hospital, Paris, France

Received 11 August 2008; revised 30 October 2008; accepted 9 February 2009

**Background:** In view of the lack of recommendations on central venous catheter (CVC)-associated thrombosis in cancer patients, we established guidelines according to the well-standardized Standards, Options and Recommendations methodology.

**Material and methods:** A literature review (1990–2007) on CVC-associated thrombosis was carried out. The guidelines were developed on the basis of the corresponding levels of evidence derived from analysis of the 36 of 175 publications selected. They were then peer reviewed by 65 independent experts.

**Results:** For the prevention of CVC-associated thrombosis, the distal tip of the CVC should be placed at the junction between the superior cava vein and right atrium; anticoagulants are not recommended. Treatment of CVC-associated thrombosis should be based on the prolonged use of low-molecular weight heparins. Maintenance of the catheter is justified if it is mandatory, functional, in the right position, and not infected, with a favorable clinical evolution under close monitoring; anticoagulant treatment should then be continued as long as the catheter is present.

**Conclusions:** Several rigorous studies do not support the use of anticoagulants for the prevention of CVC-associated thrombosis. Treatment of CVC-associated thrombosis relies on the same principles as those applied in the treatment of established thrombosis in cancer patients.

**Key words:** cancer, catheter, clinical practice guidelines, heparin, thrombosis, vitamin K antagonists

### introduction

Long-term central venous catheters (CVCs) are commonly used in patients with cancer. Their placement may be complicated by the occurrence of CVC-associated thrombosis, defined as a mural thrombus extending from the catheter into the lumen of a vessel and leading to partial or total catheter occlusion with or without clinical symptoms. In recent reviews in cancer patients, the incidence of symptomatic and asymptomatic CVC-associated thrombosis ranged between 0% and 28% and between 12 and 66%, respectively [1–4]. CVC-associated thrombosis may result in pulmonary embolism in 10%–15% of patients and loss of central venous access in

10% of patients [2]. From an economic perspective, it also accounts for a significant increase in direct treatment-related and management costs [5].

So far, no international recommendations focusing specifically on both the prophylaxis and treatment of CVC-associated thrombosis in patients with cancer (including the role of placement techniques) have been published [6]. For this reason, but also in view of recent major publications on this topic, wide heterogeneities in clinical practices, and a likely increase in the incidence of catheter thrombosis (related to an increasing incidence of cancer and a greater use of CVC), a multidisciplinary working group was set up by the French National Federation of Cancer Centers (Fédération Nationale des Centres de Lutte Contre le Cancer) to develop national guidelines for this setting according to the well-standardized procedure of the Standards, Options and Recommendations

\*Correspondence to: Dr P. Debourdeau, Hôpital Desgenettes, 108 Boulevard Pinel, 69003 Lyon, France. Tel: +33-4-72-36-61-94; Fax: +33-4-72-36-25-26; E-mail: pdebourdeau@yahoo.fr

(SOR). Initiated in 1993, the SOR program was set up to develop guidelines for the standardization of ‘good clinical practice’ throughout the various disciplines involved in cancer care [7–10]. Its methodology is based on a literature review and a critical appraisal by a multidisciplinary working group of experts [11]. It involves the cooperation of French regional cancer centers, both public and private practice sectors, scientific societies, and the French National Cancer Institute, which has led this program since May 2008.

## materials and methods

### literature review and analysis

A literature review of the studies published between January 1999 and January 2007 was carried out using the MEDLINE database and the following subject headings: cancer, thrombosis, and catheter. A prospective follow-up of the literature on this subject (meta-analyses and prospective studies only) was continued up to January 2008. National guidelines and several Evidence-Based Medicine sites were also consulted. The literature search was limited to publications in English or in French.

Meta-analyses, systematic reviews, randomized clinical trials, or nonrandomized prospective or retrospective studies in the absence of randomized clinical trials were included in the analysis. Editorials, letters to the editor, case reports, publications without an abstract, press releases, and animal studies were excluded.

We selected studies on adults and children with solid tumors or hematologic malignancies and a CVC with or without a history of thromboembolic events since the guidelines were established for both the prevention and treatment of CVC-associated thrombosis. We also analyzed publications on the role of catheter placement in CVC-associated thrombosis. We excluded studies concerning malfunctioning of catheters not related to thrombosis; these latter events are generally due to an intraluminal thrombus without mural involvement, the formation of a fibrin sleeve around the catheter, or compression of the catheter between the medial portion of the clavicle and the anterior face of the first rib (pinch-off syndrome). Likewise, we excluded studies concerning only patients with no cancer or patients with a cancer in remission for more than 5 years, a tumor-associated thrombosis, thrombocytopenia, or a catheter infection. Studies focusing on methods of diagnosing CVC-associated thrombosis were not analyzed.

For the prevention studies, the outcomes analyzed were signs and symptoms of thrombosis, asymptomatic or symptomatic thrombosis objectively confirmed (Doppler ultrasonography, contrast venography, or scanner), pulmonary embolism, or major bleeding. For the treatment studies, the outcomes analyzed were recurrence of thrombosis, pulmonary embolism, or major bleeding [12].

### critical appraisal and data extraction

The quality of the studies was evaluated with a validated reading grid assessing their methods and clinical relevance [13]. Two reviewers (Lise Bosquet, Diana Kassab Chahmi) extracted the data in a double-blind manner. Any discrepancies between reviewers were resolved by consensus.

### consensus development

Following the selection and critical appraisal of the articles, a first version of the guidelines was established based on the conclusions, the corresponding levels of evidence, and the consistency of the data (Table 1). In the absence of any clear scientific evidence, judgment was based on the professional experience and consensus of the expert group (expert agreement). In these guidelines, the recommendations were classified as Standards or Options (Table 2). The document was then peer reviewed in November 2007 by 65

**Table 1.** Definition of levels of evidence

| Level   | Definition  |
|---------|---|
| Level A | Based on one or several high-quality meta-analyses or on several high-quality randomized clinical trials with consistent results                          |
| Level B | Based on good quality evidence from randomized trials (B1) or prospective or retrospective studies (B2), with consistent results when considered together |
| Level C | Based on studies that are weak, with inconsistent results when considered together  |
| Level D | Absence of any scientific data or only a series of cases available  |

**Table 2.** Classification of recommendations

| Recommendations | Definition   |
|-----------------|--|
| Standards       | Procedures or treatments considered to be the ‘gold standard’ by unanimous decision of the experts                         |
| Options         | Procedures or treatments acknowledged to be appropriate by the experts; one of the options may be preferred by the experts |

independent experts encompassing all the medical and surgical specialties involved in the management of patients with cancer [including oncologists (33%), anesthesiologists and surgeons (9%), and hematologists (5%)] according to the AGREE grid [11], and their comments were integrated in the final version in February 2008.

## results

### primary prevention of CVC-associated thrombosis in patients with cancer

*literature search results.* Out of 175 publications on CVC-associated thrombosis, 31 publications on the primary prevention of this event in patients with cancer were identified and used for developing these guidelines [14–44].

*efficacy and safety of vitamin K antagonists.* Five randomized studies [14–18] investigated the efficacy and safety of vitamin K antagonists (VKA) in the prevention of CVC-associated thrombosis in patients with cancer (Table 3). In four studies [14–17], warfarin was administered at the once-daily dose of 1 mg/day without laboratory monitoring. In two studies, warfarin was given in order to achieve an INR (international normalized ratio) between 1.3 and 1.9 [17] or 1.5 and 2 [18]. Only the oldest study found a significant effect of VKA, compared with no treatment, in preventing CVC-associated thrombosis; this effect was obtained without increasing the risk of major bleeding [14]. VKA were not significantly more effective than placebo or no treatment in the four other later studies [15–18]. Interestingly, the percentage of CVC-associated thrombosis was lower in patients in whom warfarin was administered with a target INR between 1.5 and 2.0 than in

**Table 3.** Vitamin K antagonists in the primary prevention of CVC-associated thrombosis in patients with cancer: randomized studies from 1990 to 2007

| Reference; study date             | Type of patients; type of catheter   | No. of patients recruited                 | Treatment  | Catheter flushing                                   | Follow-up                    | End points  | Results  |
|-----------------------------------|--|---|--|---|------------------------------|---|--|
| Bern et al. [14]; not specified   | Solid tumors and lymphoma; CVC in subclavian vein (Port-a-cath)                              | 121 patients (82 patients analyzed)       | A: warfarin (1 mg/day) for 90 days;<br>B: no treatment   | UFH (up to 500 U/week)                              | 90 days                      | Asymptomatic CVC-associated DVT (venography)  | A: 9.5% (9/42); B: 37.5% (15/40), $P < 0.001$  |
| Heaton et al. [15]; not specified | Hematologic malignancies; CVC in subclavian vein (double-lumen CVC)                          | 88 patients (102 CVC but 88 CVC analyzed) | A: warfarin (1 mg/day);<br>B: no treatment   | Hickman®: UFH (50 µg, 2×/day);<br>Groshong®: saline | 90 days                      | Symptomatic thrombosis (confirmed by venography)  | A: 17.8% (8/45, with two intraluminal thromboses);<br>B: 11.6% (5/43, with one CVC-associated DVT and four intraluminal thromboses), $P = NS$                                |
| Couban et al. [16]; 1999–2002     | Solid tumors (20%), hematologic malignancies (80%); CVC (tunneled and implanted)             | 255 patients; 255 CVC                     | A: warfarin (1 mg/day) for 8 weeks (median);<br>B: placebo for 9 weeks (median)                | Not specified                                       | 25 weeks (range 1–184 weeks) | 1. Symptomatic, CVC-associated thrombosis during CVC life span (confirmed by US or venography)<br>2. Death<br>3. Major bleeding                             | 1. A: 4.6% (6/130), B: 4.0% (5/125), $P = NS$<br>2. A: 17% (22/130), B: 17% (21/125), $P = NS$<br>3. A: 0% (0/130), B: 2% (3/125), $P = NS$                                  |
| Young et al. [17]; not specified  | Solid tumors (52% colorectal cancer), hematologic malignancies; CVC (location not specified) | 1589 patients (90% analyzed)              | A: warfarin (1 mg/day) for 8 weeks (median);<br>B: warfarin (INR: 1.5–2.0);<br>C: no treatment | Not specified                                       | Not specified                | 1. Symptomatic, CVC-associated thrombosis (radiologically proven)<br>2. Major bleeding  | 1. A: 7%, B: 3%, A + B: 5%, C: 6%, A versus B: $P < 0.01$ , A + B versus C: $P = NS$<br>2. A: 2%, B: 4%, A + B: 2%, C: 0.2%, A versus B: $P = NS$ , A + B versus C: $P = NS$ |
| Ruud et al. [18]; 2002–2003       | Cancer (children); CVC in jugular vein   | 73 patients (62 patients analyzed)        | A: warfarin (INR: 1.3–1.9);<br>B: no treatment   | Not specified                                       | Not specified                | 1. Asymptomatic, CVC-associated jugular thrombosis (US at 1, 3, and 6 months)<br>2. Symptomatic DVT (CVC-associated thrombosis and PE)<br>3. Major bleeding | 1. A: 48%, B: 36%, $P = NS$<br>2. A: one symptomatic DVT, B: one symptomatic DVT<br>3. A: two major bleeds, B: zero major bleed  |

CVC, central venous catheter; DVT, deep vein thrombosis; INR, international normalized ratio; NS, not significant; PE, pulmonary embolism; UFH, unfractionated heparin; US, ultrasonography; VKA, vitamin K antagonists.

patients with a fixed dose of warfarin (3% versus 7%, respectively,  $P < 0.01$ ); however, this was obtained at the expense of a nonsignificant increase in major bleeding (4% versus 2%, respectively) [17].

Five meta-analyses evaluated the efficacy and safety of VKA in the prevention of CVC-associated thrombosis [19–23] (Table 4). None showed that VKA (either at a fixed low dose or with a target INR between 1.5 and 2.0) exerted a beneficial effect on the occurrence of symptomatic thromboses versus placebo or no treatment. However, in one meta-analysis [21], fixed low doses of VKA were more effective than placebo in preventing both asymptomatic and symptomatic CVC-associated thrombosis [relative risk = 0.37 (95% confidence interval 0.26–0.52),  $P < 0.001$ ]. Of note, this meta-analysis was not specific to cancer patients. Furthermore, these meta-analyses included a number of nonrandomized studies.

In view of the known interaction between VKA and 5-fluorouracil (5-FU), two retrospective studies (in 95 and 72 patients, respectively) [24, 25] and one noncomparative prospective study (in 247 patients with gastrointestinal cancer) [26] analyzed the effect of warfarin (1 mg/day) in cancer patients with a CVC receiving this cytotoxic drug. All showed an INR increase  $>1.5$  in 5-FU-treated patients; depending on the study, this increase was reported in 33%–50% of patients. In one study [24], the INR was  $>3.0$  in 19% of patients. Major hemorrhages were reported in 3.2%–8% of patients, 90% of these occurring in patients with a high INR.

In conclusion, the incidence of CVC-associated thrombosis in patients with cancer depended on the study. In the most recent studies, the rate of thrombosis was comparable with or without a prophylactic anticoagulant drug (~5% with regard to symptomatic thromboses) (level of evidence: A). Fixed low doses of VKA (1 mg/day) with an INR  $<1.5$  were not effective in preventing venous thrombosis associated with a superior vena cava catheter in patients with cancer (level of evidence: B1). Published data showed that the combination of low-dose VKA with 5-FU may be harmful (INR increase with consequent bleeding risk) (level of evidence: B2).

*efficacy and safety of unfractionated heparin.* Only one randomized study evaluated the efficacy and safety of unfractionated heparin (UFH) in the prevention of CVC-associated thrombosis in 108 patients with hematologic diseases (including 34 with non-malignant diseases) [27]. Patients (aged from 4 to 60 years) were randomly assigned to receive either UFH (100 U/kg/day,  $n = 65$ ) or saline ( $n = 63$ ) by continuous i.v. infusion. The CVC were externalized, nontunneled, double-lumen catheters. CVC-related asymptomatic thrombosis occurred in 1.5% of the patients treated with heparin and 12.6% of the control patients ( $P = 0.03$ ). Severe bleeding was reported in two and three patients, respectively, in the heparin and control groups ( $P = 0.18$ ).

Due to the limited number of patients and their clinical specificity (bone marrow transplantation), it was not possible to conclude on the efficacy and safety of UFH in the primary prevention of CVC-associated thrombosis in patients with cancer (level of evidence: nonevaluable).

*efficacy and safety of low-molecular weight heparins.* Six randomized studies assessing the value of low-molecular weight heparins (LMWH) in the prevention of CVC-associated thrombosis were analyzed (Table 5) [28–33]. Subcutaneous dalteparin (2500 or 5000 IU/day) was used in three studies, nadroparin (2850 IU/day) in two studies, and enoxaparin (40 mg/day) in one study.

In five studies, the comparator was either placebo [29, 30, 33] or no treatment [28, 32]. In these last two studies [28, 32], LMWH were significantly more effective in preventing asymptomatic CVC-associated thrombosis than no treatment. However, the beneficial preventive effect of LMWH, in terms of either asymptomatic or symptomatic thromboses, was not demonstrated in the three large placebo-controlled studies [30, 31, 33]. In no study was LMWH administration associated with a significant increase in the risk of bleeding. Overall, the various meta-analyses confirmed these results (Table 4). Of note, a meta-analysis combining seven studies comparing VKA, UFH, or LMWH versus placebo or no treatment in cancer patients with CVC showed that the risk of symptomatic deep vein thrombosis was significantly reduced by 44% in the group of anticoagulated patients [relative risk = 0.56 (95% confidence interval 0.34–0.92)]; there was no significant difference in the incidence of major bleeding between the two groups [22].

LMWH (dalteparin and nadroparin) were compared with fixed low-dose VKA in two studies [29, 32]. Neither study showed a statistically significant difference between the two classes of drugs in either hemorrhagic safety or efficacy. However, a meta-analysis of these two studies showed that LMWH were less effective than VKA in preventing both asymptomatic and symptomatic CVC-associated deep vein thrombosis [relative risk = 1.88 (95% confidence interval 1.28–2.75)].

In conclusion, on the basis of five concordant randomized trials of good methodological quality in patients with cancer, LMWH did not show any benefit in preventing symptomatic thromboses of the superior vena cava veins; however, they did not increase the bleeding risk (level of evidence: A) [21].

*efficacy and safety of thrombolytic drugs.* Only one nonrandomized prospective study investigated the efficacy and safety of thrombolytic drugs in the prevention of CVC-associated thrombosis [34]. This study evaluated the effect of urokinase (10 000 IU in each catheter lumen for 4 h once a week) in 15 children (16 CVC) with malignant disease; the results were compared with those obtained in a historical series of 15 children (19 CVC) without thromboprophylaxis. On systematic ultrasonography, the rate of asymptomatic thrombosis was significantly lower in the urokinase group (44%, 7 of 16 cases) than in the control group (82%, 9 of 11 cases) ( $P = 0.047$ ). No hemorrhagic complications were reported.

In view of the limited number of patients, it was not possible to conclude on the efficacy and safety of thrombolytic drugs in the primary prevention of CVC-associated thrombosis in patients with cancer (level of evidence: nonevaluable).

*influence of type, position, and method of insertion of the catheter.* A number of factors may influence the occurrence of thrombosis in patients with CVC, including the type of catheter (open-ended, such as the Hickman® catheter, versus

**Table 4.** Anticoagulant drugs in the primary prevention of CVC-associated thrombosis: meta-analyses from 1990 to 2007

| Reference                    | Number of studies analyzed; period of study selection      | No. of patients; treatment   | Thrombosis   | Other outcomes  |
|------------------------------|--|--|--|---|
| Carrier et al. [19]          | Seven studies; 1950–2007                                   | 2131 patients; VKA (warfarin 1 mg) or LMWH   | Symptomatic CVC-related thrombosis (defined as upper extremity DVT or CVC occlusion);<br>VKAs versus control: RR (95% CI) = 0.82 (0.46–1.47);<br>LMWH versus control: RR (95% CI) = 0.43 (0.12–1.56);<br>VKAs or LMWH versus control: RR (95% CI) = 0.71 (0.42–1.20)   | –   |
| Rawson and Newburn-Cook [20] | Four studies; 1966–2007                                    | 1236 patients; VKA (warfarin 1 mg or INR > 1.5)  | CVC-related thrombosis (symptomatic and asymptomatic);<br>VKAs versus control: risk difference (95% CI) = 2.0% (–9.0% to +5.0%)  | –   |
| Kirkpatrick et al. [21]      | 15 studies (10 studies on only cancer patients); 1964–2006 | 1714 patients; VKA (fixed low dose) or LMWH  | CVC-related DVT (symptomatic and asymptomatic);<br>VKAs versus control: RR (95% CI) = 0.37 (0.26–0.52), $P < 0.001$ ;<br>LMWH versus control: RR (95% CI) = 0.72 (0.57–0.90), $P = 0.045$ ;<br>LMWH versus VKAs: RR (95% CI) = 1.88 (1.28–2.75)<br>CVC-related DVT (symptomatic);<br>VKAs versus control: RR (95% CI) = 0.60 (0.30–1.20);<br>LMWH versus control: RR (95% CI) = 0.69 (0.30–1.59)   | Major bleeding:<br>VKAs versus control:<br>RR (95% CI) = 0.24 (0.03–2.13);<br>LMWH versus control:<br>RR (95% CI) = 0.66 (0.12–3.68)  |
| Akl et al. [22]              | Nine studies; from 1966                                    | 852 patients for asymptomatic DVT and 1859 patients for symptomatic DVT; VKAs or heparin (UFH or LMWH) | Asymptomatic DVT:<br>VKAs versus control: RR (95% CI) = 0.56 (0.10–2.99);<br>LMWH versus control: RR (95% CI) = 0.84 (0.52–1.36);<br>Heparin versus control: RR (95% CI) = 0.82 (0.51–1.32);<br>VKAs or heparin versus control: RR (95% CI) = 0.82 (0.73–1.68)<br>Symptomatic DVT:<br>VKAs versus control: RR (95% CI) = 0.62 (0.30–1.27);<br>LMWH versus control: RR (95% CI) = 0.49 [0.17–1.39] -<br>Heparin vs control: RR (95% CI) = 0.43 (0.18–1.06);<br>VKAs or heparin versus control: RR (95% CI) = 0.56 (0.34–0.92); $P < 0.05$ | Major bleeding: Heparin versus control:<br>RR (95% CI) = 0.68 (0.10–4.78);<br>VKAs or heparin versus control:<br>RR (95% CI) = 1.83 (0.34–9.87)<br>Death:<br>LMWH versus control: RR (95% CI) = 0.73 (0.39–1.36);<br>heparin versus control: RR (95% CI) = 0.74 (0.40–1.36);<br>VKAs or heparin versus control:<br>RR (95% CI) = 0.74 (0.40–1.36) |
| Chaukiyal et al. [23]        | Eight studies; 1966–2006                                   | 1428 patients; VKAs (warfarin 1 mg) or heparin (UFH or LMWH)   | CVC-related thrombosis (symptomatic and asymptomatic):<br>VKAs versus control: RR (95% CI) = 0.75 (0.24–2.35);<br>Heparin versus control: RR (95% CI) = 0.46 (0.18–1.20);<br>$P = 0.06$ ;<br>VKAs or heparin versus control: RR (95% CI) = 0.59 (0.31–1.13); $P = 0.11$ ;<br>VKAs versus LMWH: RR (95% CI) = 1.71 (0.56–5.26)  | Major bleeding:<br>VKAs versus control:<br>RR (95% CI) = 0.14 (0.01–2.63);<br>heparin versus control:<br>RR (95% CI) = 0.41 (0.05–3.30);<br>VKAs or heparin versus control:<br>RR (95% CI) = 0.44 (0.12–1.67)   |

CI, confidence interval; CVC, central venous catheter; DVT, deep vein thrombosis; INR, international normalized ratio; LMWH, low-molecular weight heparins; RR, relative risk; UFH, unfractionated heparin; VKAs: vitamin K antagonists.

**Table 5.** Low-molecular weight heparins in the primary prevention of CVC-associated thrombosis in patients with cancer: randomized studies from 1990 to 2007

| Reference; study date              | Type of patients; type of catheter   | No. of patients                      | Treatment   | Catheter flushing                                   | Follow-up     | End points   | Results   |
|------------------------------------|--|--------------------------------------|---|---|---------------|--|---|
| Monreal et al. [28]; 1993–1995     | Solid tumors; CVC (Port-a-cath)  | 32 patients; (29 patients analyzed)  | A: dalteparin (2500 IU/day);<br>B: no treatment   | Heparinized saline (10 ml, once a week)             | 90 days       | 1. Asymptomatic CVC-associated DVT (venography)<br>2. Major bleeding   | 1. A: 6.2% (1/16), B: 61.5% (8/13), $P = 0.002$<br>2. A: one patient, B: zero patient   |
| Mismetti et al. [29]; 1998–2000    | Solid tumors; CVC (totally implantable port-system CVC) in subclavian (95%) or jugular vein  | 59 patients (45 patients analyzed)   | A: nadroparin (2850 IU/day) for 90 days;<br>B: warfarin (1 mg/day) for 90 days  | Saline (10 ml) and heparinized saline (500 U, 5 ml) | 6 months      | 1. Asymptomatic and symptomatic CVC-associated DVT (venography) at 90 days<br>2. All VTE events at 90 days<br>3. All VTE events at 6 months<br>4. Major bleeding | 1. A: 28.6% (6/21), B: 16.7% (4/24), $P = NS$<br>2. A: 31.8%, B: 16.7%, $P = NS$<br>3. A: 36.4%, B: 16.7%, $P = NS$<br>4. A: one patient, B: zero patient |
| Verso et al. [30]; 2000–2003       | Solid tumors (54% of gastrointestinal cancer), hematologic malignancies (9%); CVC (polyurethane or silicone) in subclavian (89%) or other vein | 385 patients (310 patients analyzed) | A: enoxaparin (40 mg/day) for 6 weeks;<br>B: placebo for 6 weeks  | Not specified                                       | 3 months      | 1. Composite of asymptomatic or symptomatic upper limb DVT (venography) or symptomatic PE<br>2. Major bleeding<br>3. Death                                       | 1. A: 14.1% (22/155), B: 18.0% (28/155), $P = NS$<br>2. A: 0, B: 0<br>3. A: 2.6% (5/191), B: 1.0% (1/194)   |
| Karthaus et al. [31]; 1999–2001    | Solid tumors (90%), hematologic malignancies; CVC  | 439 patients (425 patients analyzed) | A: dalteparin (5000 IU/day) for 16 weeks;<br>B: placebo for 16 weeks  | UFH (500 U) in saline solution during CVC use       | 16 weeks      | 1. Symptomatic CVC-associated DVT (venography, US, CT scan)<br>2. Asymptomatic CVC-associated DVT (venography, US)<br>3. Bleeding events                         | 1. A: 3.7% (11/294), B: 3.4% (5/145), $P = NS$<br>2. A: 3.4% (10/294), B: 4.1% (6/145), $P = NS$<br>3. A: 17.5% (50/285), B: 15.0% (21/140), $P = NS$     |
| DeCicco et al. [32]; not specified | Cancer; CVC  | 450 patients (348 patients analyzed) | A: acenocoumarin (1 mg/day) initiated 3 days before CVC insertion and continued for 8 days after CVC insertion;<br>B: dalteparin (5000 IU/day) for 8 days after CVC insertion;<br>C: no treatment | Not specified                                       | Not specified | 1. Asymptomatic CVC-associated DVT (venography)<br>2. PE<br>3. Major bleeding  | 1. A: 21.9%, B: 40%, C: 55.3%; A versus B: $P = 0.003$ ; A versus C: $P < 0.001$ ; B versus C: $P < 0.02$<br>2. No PE<br>3. No major bleeding             |
| Niers et al. [33]; not specified   | Hematologic malignancies; CVC (chemotherapy and stem-cell transplantation)   | 113 patients (87 patients analyzed)  | A: nadroparin (2850 IU/day) for 3 weeks;<br>B: placebo for 3 weeks  | Not specified                                       | 3 weeks       | 1. Asymptomatic CVC-associated DVT (venography)<br>2. Major bleeding   | 1. A: 17% (7/41), B: 9% (4/46), $P = 0.49$<br>2. A: 0, B: 0   |

CVC, central venous catheter; DVT, deep-vein thrombosis; NS, not significant; PE, pulmonary embolism; UFH, unfractionated heparin; US, ultrasonography; VTE, venous thromboembolism.

closed-ended catheter with a valve, such as the Groshong® catheter), its position (above, below, or at the junction of the superior cava vein and the right atrium), and the method of placement. The analysis of the role of these factors in CVC-associated thrombosis was based on two randomized studies, five nonrandomized prospective studies, and four retrospective series (Tables 6–8) [35–44].

Closed-ended or valved catheters were compared with open-ended or nonvalved catheters in two randomized studies (Table 6) [35, 36]. None of these studies showed any significant difference between the two study groups in terms of symptomatic thrombosis.

The influence of the position of the catheter tip on CVC-associated thrombosis was assessed in six nonrandomized studies: in four of these studies [38, 41, 43, 44], a higher rate of thrombosis was observed when the CVC tip was located above the junction between the superior cava vein and the right atrium. Three studies also reported that a left-sided insertion of CVC significantly increased the risk of thrombotic complications [39, 42, 43]. Other risk factors for symptomatic thrombosis were femoral position of the CVC, a duration of placement >25 min [39], and more than one CVC placement attempt [40].

In conclusion, the concordant data of these studies highlighted the lower thrombogenicity of some placement characteristics of CVC, i.e. (i) distal tip at the junction of the superior cava vein and the right atrium (level of evidence: B2) and (ii) whenever possible, right-sided insertion (level of evidence: B2). Conversely, various placement characteristics increase the risk of CVC-associated thrombosis, i.e. (i) number of attempts (more than two) and duration of placement (level of evidence: D) and (ii) placement of the CVC in a femoral vein (level of evidence: D).

### treatment of CVC-associated thrombosis in patients with cancer

*literature search results.* Out of 175 publications on CVC-associated thrombosis, five publications on the curative treatment of CVC-associated thrombosis in patients with cancer were identified and used for developing these guidelines [45–49].

*efficacy and safety of LMWH.* Only one nonrandomized prospective study examined the efficacy and safety of LMWH in the treatment of CVC-associated thrombosis [45]. In this study, 46 outpatients (34 with a cancer and 16 with a CVC) with confirmed upper extremity deep vein thrombosis were treated with dalteparin (200 IU/kg once daily for a minimum of 5 days) followed by warfarin (target INR: 2.0–3.0); at 12 weeks, there was one recurrence of deep vein thrombosis confirmed by Doppler ultrasonography or venography and one major bleeding.

Based on the published data (only one low-quality study), it was not possible to conclude on the efficacy and safety of short-term LMWH followed by VKA in the curative treatment of CVC-associated thrombosis in patients with cancer (level of evidence: nonevaluable).

However, the experts proposed that, on the basis of good quality studies showing concordant results concerning the efficacy and safety of LMWH given for 3–6 months in the treatment of deep vein thrombosis of lower limbs or pulmonary

**Table 6.** Influence of type, position, and method of insertion of catheter in the primary prevention of CVC-associated thrombosis: randomized studies from 1990 to 2007

| Reference; study date            | Type of study; type of patients | No. of patients recruited | Treatment   | Catheter flushing  | Follow-up                     | End points  | Results   |
|----------------------------------|---------------------------------|---------------------------|---|--|-------------------------------|---|---|
| Biffi et al. [35]; 1997–1998     | Randomized study; solid tumors  | 304 patients (302 CVC)    | A: 8.0-F, silastic Groshong® CVC;<br>B: 9.6-F, silastic, open-ended CVC | Saline (20 ml), then heparinized saline after each CVC use (5 ml at 50 U/ml) | 237 days                      | 1. Asymptomatic and asymptomatic DVT in internal jugular or subclavian vein (venography)<br>2. CVC removal<br>3. Clinically relevant bleeding | 1. A: 3.9% (6/152), B: 7.3% (11/150), P = NS<br>2. A: 2.7% (4/152), B: 3.3% (5/150)<br>3. No clinically relevant bleeding |
| Carlo et al. [36]; not specified | Randomized study; solid tumors  | 73 patients               | A: valved port;<br>B: nonvalved port                                    | A: saline (10 ml);<br>B: heparinized saline (10 ml)                          | 180 days or until CVC removal | Symptomatic thrombosis of internal jugular vein   | A: 2.7% (1/37), B: 2.8% (1/36), P = NS  |

CVC, central-venous catheter; DVT, deep vein thrombosis; NS, not significant.

**Table 7.** Influence of type, position, and method of insertion of catheter in the primary prevention of CVC-associated thrombosis: prospective studies from 1990 to 2007

| Reference; study date              | Type of study; type of patients   | No. of patients recruited             | Treatment           | Catheter flushing   | Follow-up         | End points   | Results  |
|------------------------------------|---|---------------------------------------|---------------------|---|-------------------|--|--|
| Nightingale et al. [37]; 1993–1994 | Nonrandomized prospective comparative study; gastrointestinal cancer; tunneled CVC in right subclavian (727), left subclavian (81), right femoral (two) or jugular (one) vein | 949 patients; (832 patients analyzed) | Warfarin (1 mg/day) | Heparinized saline  | Not specified     | 1. Thrombotic complications leading to CVC removal<br>2. Predictive factor for CVC removal (multivariate analysis)                   | 1. 4.7% (38/817); if distal CVC tip in SVC: 3.5% (20/569);<br>if distal CVC tip in right atrium: 2.5%, (4/160), <i>P</i> = NS<br>2. CVC in SVC: HR ([95% CI] = 2.57)   |
| Luciani et al. [38]; 1995–1998     | Nonrandomized prospective comparative study; oropharyngeal tract cancer; totally implantable CVC  | 145 patients (113 CVC)                | Not specified       | Saline (10 ml), then heparinized saline (5 ml at 50 U/ml) | >3 years          | Asymptomatic and symptomatic CVC-associated DVT (Doppler US)   | 11.7% (17/145), 76% of which were asymptomatic<br>1. If distal CVC tip in the SVC or at the junction between SVC and right atrium: 6% (5/87); if distal CVC tip above the junction between SVC and right atrium: 46% (12/26), <i>P</i> < 0.001<br>2. Left-sided CVC: 65% (11/17); right-sided CVC: 35% (6/17), <i>P</i> = NS |
| Morazin et al. [39]; 1995–1999     | Nonrandomized prospective comparative study; solid tumors (50% breast cancer); tunneled CVC (silicone)  | 5447 CVC                              | Not specified       | Not specified   | Up to CVC removal | Predictive factors for symptomatic CVC-associated DVT (venography, Doppler US, contrast computed tomography) (multivariate analysis) | 2.5% (135/5447)<br>1. Left subclavian vein + jugular vein versus right subclavian vein: RR = 2.6, <i>P</i> < 0.001<br>2. Femoral vein versus right subclavian vein: RR = 6.5, <i>P</i> < 0.001<br>3. Placement duration >25 min versus ≤25 min: RR = 1.52, <i>P</i> = 0.02   |



Table 7. (Continued)

| Reference; study date      | Type of study; type of patients   | No. of patients recruited | Treatment     | Catheter flushing   | Follow-up   | End points  | Results  |
|----------------------------|---|---------------------------|---------------|---|---|---|--|
| Lee et al. [40]; 2002–2003 | Nonrandomized prospective comparative study; solid tumors (66%); hematologic malignancies (34%); all types of CVC in the upper limb vasculature | 444 patients (555 CVC)    | Not specified | Implanted ports: heparinized saline (100 U/ml); other CVC: saline | Up to CVC removal + 4 weeks or up to 52 weeks after CVC insertion | Predictive factors for symptomatic CVC-associated DVT (US, venography, contrast computed tomography, or magnetic resonance imaging) (multivariate analysis) | 1. >1 insertion attempts: OR (95% CI) = 5.5 (1.2–24.6), $P = 0.03$<br>2. Previous CVC insertion: OR (95% CI) = 3.8 (1.4–10.4), $P = 0.01$<br>3. CVC blockage: OR (95% CI) = 14.7 (5.5–40), $P < 0.001$ |

CI, confidence interval; CVC, central venous catheter; DVT, deep vein thrombosis; HR, hazard ratio; NS, not significant; OR, odds ratio; RR, relative risk; SVC, superior vena cava; US, ultrasonography.

embolism in patients with cancer [50], the prolonged use of LMWH alone may be considered for the treatment of CVC-associated thrombosis, depending on the clinical status of the patient. By analogy with the management of patients with venous thromboembolism and renal insufficiency [51], the experts recommended that in the event of severe renal impairment, the treatment should be based on the use of UFH, rapidly followed (possibly as early as the first day) by VKA.

*efficacy and safety of thrombolytic drugs.* The value of thrombolytic drugs in the treatment of CVC-associated thrombosis was assessed in two nonrandomized prospective studies [46, 47] and one retrospective study [48], each with a limited number of patients. In the first, small study, only four adults and one child with cancer and CVC-associated thrombosis were treated with a continuous infusion of both recombinant tissue-type plasminogen activator (0.5 mg/kg per 24 h, preceded by a 5-mg bolus injection in adult patients or a 2-mg bolus injection in the child) and UFH for 4.5–7.9 days [46]. The treatment was effective in resolving large vessel obstruction without bleeding in three out of five patients. Partial lysis of the thrombus and moderately severe hemorrhage were observed in the other two patients.

The second study concerned 18 cancer patients receiving high-dose chemotherapy who developed CVC-associated thrombosis [47]. These patients were treated with urokinase (750 000–150 000 U/h for 24–96 h) infused into a vein of the ipsilateral upper limb. A partial or complete resolution of clinical signs and symptoms was reported in all patients. A partial radiographic response was found in nine patients (50%). Major bleeding was observed in four patients.

The third study was a retrospective comparison of the efficacy of various thrombolytic drugs versus LMWH in 57 patients with CVC-associated thrombosis [48]. Thirty-two patients received a thrombolytic drug, streptokinase ( $n = 16$ ), urokinase ( $n = 5$ ), tissue plasminogen activator ( $n = 4$ ), or a combination of streptokinase and urokinase ( $n = 7$ ), via a systematic route. Repermeabilization (as assessed by systematic Doppler ultrasonography) was observed in 16 patients (50%). No serious side-effects were observed. By comparison, in 25 patients treated with curative doses of enoxaparin for 3 weeks followed by warfarin, repermeabilization was observed in only one patient (5%,  $P = 0.009$  versus thrombolytic drugs).

In conclusion, it was not possible to conclude on the efficacy and safety of thrombolytic drugs, administered either systemically or locally. Published data have shown the feasibility of their administration, including in patients treated with intensive chemotherapy (level of evidence: D).

Thus, the experts proposed that, based on published data, the administration of thrombolytic drugs for the treatment of CVC-associated thrombosis may only be considered in specific circumstances, in which the thrombotic risk is superior to the risk associated with the use of these drugs, i.e. in the event of superior vena cava thrombosis associated with recent, poorly tolerated, vena cava syndrome objectively confirmed (at least on a thoracic computed tomography scan and/or opacification of the superior vena cava), or imperative maintenance of a CVC.

**Table 8.** Influence of type, position, and method of insertion of catheter in the primary prevention of CVC-associated thrombosis: retrospective studies from 1990 to 2007

| Reference; study date               | Type of study; type of patients  | No. of patients recruited  | Treatment     | Catheter flushing  | Follow-up             | End points   | Results   |
|-------------------------------------|--|--|---------------|--|-----------------------|--|---|
| Eastridge and Lefor [41]; 1989–1992 | Nonrandomized retrospective comparative study; solid tumors (51%); hematologic malignancies (49%); tunneled CVC (65%); implantable CVC (35%) | 274 patients (332 CVC)   | Not specified | Heparinized saline (3 ml/day at 100 U/ml)                  | Not specified         | Predictive factors for symptomatic CVC-associated DVT (venography or clinical follow-up)               | <ol style="list-style-type: none"> <li>Position of CVC tip: above third dorsal vertebra = 78% of patients with CVC-associated thrombosis (versus 37% if below third dorsal vertebra), <math>P &lt; 0.05</math></li> <li>Triple lumen CVC = 21% (10/48); double lumen CVC = 7% (11/160), <math>P &lt; 0.05</math></li> <li>Implantable CVC = 6% (7/113); tunneled CVC = 10% (21/209), <math>P = NS</math></li> </ol> |
| Craft et al. [42]; not specified    | Nonrandomized retrospective comparative study; solid tumors (48%); hematologic malignancies (41%); tunneled CVC (Hickman®)                   | 122 patients (120 patients analyzed); 153 CVC (150 CVC analyzed) | Not specified | Heparinized saline   | 55 days (range 1–650) | Predictive factors for symptomatic CVC-associated DVT (venography)                                     | <ol style="list-style-type: none"> <li>Position of CVC tip: junction right atrium SVC or lower third of SVC: 8.2%; upper third of SVC: 7.5%, <math>P = NS</math></li> <li>Side of CVC: right sided: 5%; left sided: 19%, RR (95% CI) = 4.4 (1.2–16), <math>P = 0.04</math></li> </ol>   |
| Cadman et al. [43]; 1996–2001       | Randomly sampled retrospective study; solid tumors (69%); hematologic malignancies (31%); tunneled CVC                                       | 334 patients (448 CVC)   | Not specified | Not specified  | 72 days (range 1–720) | Predictive factors for symptomatic CVC-associated DVT (venography, Doppler US)                         | <ol style="list-style-type: none"> <li>Position of CVC tip: right atrium: 0%; lower third of SVC: 2.6%; middle third of SVC: 5.3%; upper third of SVC: 41.7%, <math>P &lt; 0.005</math></li> <li>Side of CVC placement: right sided: 6.8%; left sided: 25.6%, <math>P &lt; 0.001</math></li> </ol>  |
| Caers et al. [44]; 1993–1998        | Nonrandomized retrospective comparative study; solid tumors (84%); hematologic malignancies (13%)  | 437 patients (448 CVC)   | Not specified | Saline (10 ml), then heparinized saline (5 ml at 100 U/ml) | –                     | Predictive factors for symptomatic CVC-associated DVT (venography, Doppler US) (multivariate analysis) | <ol style="list-style-type: none"> <li>CVC tip in the brachiocephalic vein: OR (95% CI) = 64.7 (7.6–553.8)</li> <li>CVC tip in the cranial part of the SVC: OR (95% CI) = 17.4 (2.0–148.8)</li> </ol>   |

CI, confidence interval; CVC, central venous catheter; DVT, deep vein thrombosis; NS, not significant; OR, odds ratio; RR, relative risk; SVC, superior vena cava; US, ultrasonography.

*evaluation of catheter removal.* One retrospective study evaluated the benefit of CVC removal in patients with CVC-associated thrombosis [49]. In this study, in 319 cancer patients, 112 (35%) exhibited CVC-associated thrombosis on radionuclide venography. Various therapeutic interventions, including anticoagulation with heparin or warfarin or both; line removal or replacement; or a combination thereof, were carried out. Overall, the catheter was removed in 52% of these patients. Only four patients did not show resolution of their presenting symptoms; they were treated by line replacement. No patient experienced pulmonary embolism.

In conclusion, the published data are insufficient (only one retrospective study with methodological biases) to conclude on the value of catheter removal. In the event of catheter removal, no data were reported on the optimal interval between removal and initiation of anticoagulant treatment (level of evidence: nonevaluable).

The experts did not recommend catheter removal if all the following conditions are met: (i) the distal catheter tip is in the right position (at the junction between the superior vena cava and the right atrium), (ii) the catheter is functional (good blood reflux), (iii) the catheter is mandatory or vital for the patient, and (iv) there is no fever or any sign or symptom of infected thrombophlebitis. In contrast, catheter removal is warranted if there is a prime risk factor for thrombosis (catheter too short, misplaced, etc.). There are no reliable data on the optimal duration of anticoagulant treatment after catheter removal.

## conclusion

Based on the literature review and the well-argued judgment of experts, the 2008 SOR guidelines for the prevention and treatment of CVC-associated thrombosis in patients with cancer are as follows.

### primary prevention of CVC-associated thrombosis in patients with cancer

*standards.*

- 1 The distal tip of CVC should be placed at the junction between the superior vena cava and the right atrium.
- 2 The primary prevention of CVC-associated thrombosis with anticoagulant drugs is not recommended in patients with cancer.

*options.*

- 1 Right-sided insertion and placement of the CVC in a specialized unit should be favored.

### treatment of CVC-associated thrombosis in patients with cancer

*standards.*

- 1 The treatment of CVC-associated thrombosis should be based on the prolonged use of LMWH.
- 2 In the event of severe renal impairment, the treatment should be based on the use of UFH, rapidly followed (possibly as early as the first day) by VKA.

- 3 Maintenance of the catheter is justified in the event that the catheter is mandatory, functional, in the right position, and not infected, with a favorable clinical evolution under close monitoring. In this case, an anticoagulant treatment should be maintained as long as the catheter is present.
- 4 In the event of catheter removal, there is no standard approach in terms of the interval between removal and initiation of anticoagulant treatment.

*options.*

- 1 If it is necessary to place a new catheter, the status of the superior vena cava network should be evaluated by a scan or Doppler ultrasonography.
- 2 In the event of refusal or impossibility of a prolonged treatment with LMWH, short-term use of LMWH followed by VKA may be proposed.
- 3 Thrombolytic drugs may be considered in specialized units in the event of poor clinical tolerance (vena cava syndrome) and in the absence of any contraindications.
- 4 There are no reliable data on the optimal duration of anticoagulant treatment after catheter removal.

Of note, for all recommendations classified as 'Options', further research is warranted. Moreover, none of these recommendations applies to patients with tumor-associated thrombosis or to patients with catheters that are malfunctioning for reasons other than thrombosis (fibrin sleeve, pinch-off, intraluminal thrombus), infected catheters, or femoral catheters.

## funding

French Federation of Cancer Centers and the French National Cancer League.

## acknowledgements

The authors acknowledge the support of the French Federation of Cancer Centers, the *Ligue Nationale contre le Cancer*, the *Fédération Hospitalière de France*, the *Fédération Nationale de Cancérologie des CHRU*, the *Fédération Française de Cancérologie des CHG*, the *Union Nationale Hospitalière Privée de Cancérologie*, the *Société Française de Médecine Vasculaire*, and the *Société Nationale Française de Médecine Interne*.

The authors thank:

The other members of the SOR working group: Francis Cajfinger, Hôpital Pitié Salpêtrière, Paris; Hélène Desmurs-Clavel, Hôpital Édouard Herriot, Lyon; Antoine Elias, Hôpital Font Pré, Toulon; Claire Grange, Hôpital Lyon Sud, Lyon; Hamid Hocini, Hôpital Saint Louis, Paris; Isabelle Mahé, Hôpital Louis Mourier, Colombes.

The following reviewers: Thierry André, Hôpital Pitié Salpêtrière, Paris; Elias Assaf, Hôpital Henri Mondor, Créteil; Marie-Françoise Avril, Hôpital Cochin, Paris; Marie-Thérèse Barrelier, CHU Côte de Nacre, Caen; Jean Michel Baud, Cabinet privé, Chesnay; François Becker, Hôpital Cantonal, Genève; Jean-François Bergmann, Hôpital Lariboisière, Paris; Jean-Yves Blay, Hôpital Edouard Herriot, Lyon; Philippe

Bouju, CHI Robert Ballanger, Aulnay-sous-Bois; Luc Bressolette, Hôpital de la Cavale Blanche, Brest; Pauline Brice, Hôpital Saint-Louis, Paris; Pascal Cathebras, Hôpital Nord, Saint-Étienne; Bruno Chauffert, Centre Georges François Leclerc, Dijon; Gisèle Chvetzoff, Centre Léon Bérard, Lyon; Thierry Conroy, Centre Alexis Vautrin, Vandœuvre-Lès-Nancy; Joel Constans, Hôpital Saint André, Bordeaux; Paul Cottu, Institut Curie, Paris; Francis Couturaud, CHU de Brest, Brest; Didier Cupissol, Centre Val d'Aurelle, Montpellier; Marc Espié, Hôpital Saint-Louis, Paris; Pierre-Luc Etienne, Clinique Armoricaine de Radiologie, Saint-Brieuc; Philippe Girard, Institut Mutualiste Monsouris, Paris; Bernard Goichot, Hôpital de Haute-pierre, Strasbourg; Jean-Paul Guastalla, Centre Léon Bérard, Lyon; Daniel Hayoz, Hôpital Cantonal Fribourg, Lausanne; Jean-Léon Lagrange, Hôpital Henri Mondor, Créteil; Jean-Pierre Laroche, Cabinet Angéiologie, Avignon; Alain Le Quellec, Hôpital Saint-Eloi, Montpellier; Thomas Lecompte, CHU de Nancy, Vandoeuvre-Lès-Nancy; Claire Le Hello, CHU Côte de Nacre, Caen; Christophe Leroyer, Hôpital de la Cavale Blanche, Brest; Anne Long, Hôpital Robert Debré, Reims; Alain Lortholary, Centre Catherine de Sienne, Nantes; Nadine Magy-Bertrand, CHU Jean Minjoz, Besançon; Jean-Baptiste Meric, Hôpital Pitié Salpêtrière, Paris; Laurent Mineur, Institut Sainte Catherine, Avignon; Jane Muret, Institut Gustave Roussy, Villejuif; Dominique Musset, Hôpital Antoine Béclère, Clamart; Sylvie Négrier, Centre Léon Bérard, Lyon; Martine Pacailler, Clinique Trelen, Sainte Colombe; Erwan Papin, Cabinet privé, Saint-André-de-Cubzac; Gilles Pernod, CHU de Grenoble, Grenoble; Eric Perrier, HIA Percy, Clamart; Pierre Philippe, CHU Hôtel Dieu, Clermont-Ferrand; Pascal Piedbois, Hôpital Henri Mondor, Créteil; Laurent Pinède, Clinique Protestante, Caluire-et-Cuire; Jean-Luc Reny, CH de Béziers, Béziers; Marc Righini, Hôpital Cantonal, Genève; Pascal Roblot, CHU, Poitiers; Luc Ronchi, CH de Saint-Nazaire, Saint-Nazaire; Marc Samama, Hôpital Hôtel Dieu, Paris; François Saunier, CHU de Lyon Sud, Pierre-Bénite; Stéphane Schneider, Hôpital de l'Archet, Nice; Jean-François Schved, Hôpital Saint-Eloi, Montpellier; Marie-Antoinette Sevestre, Hôpital Sud CHU, Amiens; Pascal Sève, Hôpital Hôtel Dieu, Lyon; Annick Steib, Hôpital civil, Strasbourg; Antoine Thyss, Centre Antoine Lacassagne, Nice; Jean Tredaniel, Hôpital Saint-Louis, Paris; Bruno Tribout, Hôpital Sud CHU, Amiens; Jean-Pierre Vannier, Hôpital Charles Nicolle, Rouen; Michel Vayssairat, Hôpital Tenon, Paris; Eric Voog, Clinique Victor Hugo, Le Mans; Denis Wahl, CHU de Nancy, Nancy; Marc Ychou, Centre Val d'Aurelle, Montpellier.

## references

- Klerk CP, Smorenburg SM, Buller HR. Thrombosis prophylaxis in patient populations with a central venous catheter: a systematic review. *Arch Intern Med* 2003; 163: 1913–1921.
- Verso M, Agnelli G. Venous thromboembolism associated with long-term use of central venous catheters in cancer patients. *J Clin Oncol* 2003; 21: 3665–3675.
- Boersma RS, Jie KS, Verbon A et al. Thrombotic and infectious complications of central venous catheters in patients with hematological malignancies. *Ann Oncol* 2008; 19: 433–442.
- Vescia S, Baumgärtner AK, Jacobs VR et al. Management of venous port systems in oncology: a review of current evidence. *Ann Oncol* 2008; 19: 9–15.
- Kuter DJ. Thrombotic complications of central venous catheters in cancer patients. *Oncologist* 2004; 9: 207–216.
- Geerts WH, Pineo GF, Heit JA et al. Prevention of venous thromboembolism: the Seventh ACCP Conference on Antithrombotic and Thrombolytic Therapy. *Chest* 2004; 126 (3 Suppl): 338S–400S.
- Fervers B, Hardy J, Blanc-Vincent MP et al. SOR: project methodology. *Br J Cancer* 2001; 84 (Suppl 2): 8–16.
- Burgers JS, Fervers B, Haugh M et al. International assessment of the quality of clinical practice guidelines in oncology using the Appraisal of Guidelines and Research and Evaluation Instrument. *J Clin Oncol* 2004; 22: 2000–2007.
- Fervers B, Burgers JS, Haugh MC et al. Predictors of high quality clinical practice guidelines: examples in oncology. *Int J Qual Health Care* 2005; 17: 123–132.
- Standards, Options: Recommendations. <http://www.sor-cancer.fr/> (31 July 2008, date last accessed).
- AGREE Collaboration. Development and validation of an international appraisal instrument for assessing the quality of clinical practice guidelines: the AGREE project. *Qual Saf Health Care* 2003; 12: 18–23.
- Schulman S, Kearon C. Subcommittee on Control of Anticoagulation of the Scientific and Standardization Committee of the International Society on Thrombosis and Haemostasis. Definition of major bleeding in clinical investigations of antihemostatic medicinal products in non-surgical patients. *J Thromb Haemost* 2005; 3: 692–694.
- Cucherat M. Guide de lecture critique d'un essai thérapeutique. *Médecine Thérapeutique* 2006; 12: 354–358. <http://www.spc.univ-lyon1.fr/lecture-critique/grille%20de%20lecture/frame1.htm> (31 July 2008, date last accessed).
- Bern MM, Lokich JJ, Wallach SR et al. Very low doses of warfarin can prevent thrombosis in central venous catheters. A randomized prospective trial. *Ann Intern Med* 1990; 112: 423–428.
- Heaton DC, Han DY, Inder A. Minidose (1 mg) warfarin as prophylaxis for central vein catheter thrombosis. *Intern Med J* 2002; 32: 84–88.
- Couban S, Goodyear M, Burnell M et al. Randomized placebo-controlled study of low-dose warfarin for the prevention of central venous catheter-associated thrombosis in patients with cancer. *J Clin Oncol* 2005; 23: 4063–4069.
- Young AM, Begum J, Billingham LJ et al. WARP—a multicentre prospective randomised controlled trial (RCT) of thrombosis prophylaxis with warfarin in cancer patients with central venous catheters (CVCs). *J Clin Oncol* 2005; 16s (Suppl): (Abstr LBA8004).
- Ruud E, Holmstrom H, De Lange C et al. Low-dose warfarin for the prevention of central line-associated thromboses in children with malignancies—a randomized, controlled study. *Acta Paediatr* 2006; 95: 1053–1059.
- Carrier M, Tay J, Fergusson D, Wells PS. Thromboprophylaxis for catheter-related thrombosis in patients with cancer: a systematic review of the randomized, controlled trials. *J Thromb Haemost* 2007; 5: 2552–2554.
- Rawson KM, Newburn-Cook CV. The use of low-dose warfarin as prophylaxis for central venous catheter thrombosis in patients with cancer: a meta-analysis. *Oncol Nurs Forum* 2007; 34: 1037–1043.
- Kirkpatrick A, Rathbun S, Whitsett T, Raskob G. Prevention of central venous catheter-associated thrombosis: a meta-analysis. *Am J Med* 2007; 120: 901–913.
- Akl EA, Karmath G, Yosuico V et al. Anticoagulation for thrombosis prophylaxis in cancer patients with central venous catheters. *Cochrane Database Syst Rev* 2007; CD006468.
- Chaukiyal P, Nautiyal A, Radhakrishnan S et al. Thromboprophylaxis in cancer patients with central venous catheters. A systematic review and meta-analysis. *Thromb Haemost* 2008; 99: 38–43.
- Masci G, Magagnoli M, Zucali PA et al. Minidose warfarin prophylaxis for catheter-associated thrombosis in cancer patients: can it be safely associated with fluorouracil-based chemotherapy? *J Clin Oncol* 2003; 21: 736–739.
- Magagnoli M, Masci G, Carnaghi C et al. Minidose warfarin is associated with a high incidence of International Normalized Ratio elevation during chemotherapy with FOLFOX regimen. *Ann Oncol* 2003; 14: 959–960.
- Magagnoli M, Masci G, Castagna L et al. High incidence of INR alteration in gastrointestinal cancer patients treated with mini-dose warfarin and 5-fluorouracil-based regimens. *Ann Oncol* 2006; 17: 174–176.

27. Abdelkefi A, Ben OT, Kammoun L et al. Prevention of central venous line-related thrombosis by continuous infusion of low-dose unfractionated heparin, in patients with haemato-oncological disease. *Thromb Haemost* 2004; 92: 654–661.
28. Monreal M, Alastrue A, Rull M et al. Upper extremity deep venous thrombosis in cancer patients with venous access devices—prophylaxis with a low molecular weight heparin (Fragmin). *Thromb Haemost* 1996; 75: 251–253.
29. Mismetti P, Mille D, Laporte S et al. Low-molecular-weight heparin (nadroparin) and very low doses of warfarin in the prevention of upper extremity thrombosis in cancer patients with indwelling long-term central venous catheters: a pilot randomized trial. *Haematologica* 2003; 88: 67–73.
30. Verso M, Agnelli G, Bertoglio S et al. Enoxaparin for the prevention of venous thromboembolism associated with central vein catheter: a double-blind, placebo-controlled, randomized study in cancer patients. *J Clin Oncol* 2005; 23: 4057–4062.
31. Karthaus M, Kretzschmar A, Kroning H et al. Dalteparin for prevention of catheter-related complications in cancer patients with central venous catheters: final results of a double-blind, placebo-controlled phase III trial. *Ann Oncol* 2006; 17: 289–296.
32. DeCicco M, Matovic M, Pacenzia R et al. Short-term acenocumarine (A) or dalteparine (D) for the prevention of central venous catheter-related thrombosis (CVCrT) in cancer patients. A randomized controlled study based on serial venographies. *J Clin Oncol* 2006; 18S (Suppl): (Abstr 8549).
33. Niers TM, Di NM, Klerk CP et al. Prevention of catheter-related venous thrombosis with nadroparin in patients receiving chemotherapy for hematologic malignancies: a randomized, placebo-controlled study. *J Thromb Haemost* 2007; 5: 1878–1882.
34. Kalmanti M, Germanakis J, Stiakaki E et al. Prophylaxis with urokinase in pediatric oncology patients with central venous catheters. *Pediatr Hematol Oncol* 2002; 19: 173–179.
35. Biffi R, De Braud F, Orsi F et al. A randomized, prospective trial of central venous ports connected to standard open-ended or Groshong catheters in adult oncology patients. *Cancer* 2001; 92: 1204–1212.
36. Carlo JT, Lamont JP, McCarty TM et al. A prospective randomized trial demonstrating valved implantable ports have fewer complications and lower overall cost than nonvalved implantable ports. *Am J Surg* 2004; 188: 722–727.
37. Nightingale CE, Norman A, Cunningham D et al. A prospective analysis of 949 long-term central venous access catheters for ambulatory chemotherapy in patients with gastrointestinal malignancy. *Eur J Cancer* 1997; 33: 398–403.
38. Luciani A, Clement O, Halimi P et al. Catheter-related upper extremity deep venous thrombosis in cancer patients: a prospective study based on Doppler US. *Radiology* 2001; 220: 655–660.
39. Morazin F, Kriegel I, Asselain B, Falcou MC. Thrombose symptomatique sur cathéter veineux central de longue durée en oncologie: un score de risque prédictif? *Rev Med Intern* 2005; 26: 273–279.
40. Lee AY, Levine MN, Butler G et al. Incidence, risk factors, and outcomes of catheter-related thrombosis in adult patients with cancer. *J Clin Oncol* 2006; 24: 1404–1408.
41. Eastridge BJ, Lefor AT. Complications of indwelling venous access devices in cancer patients. *J Clin Oncol* 1995; 13: 233–238.
42. Craft PS, May J, Dorigo A et al. Hickman catheters: left-sided insertion, male gender, and obesity are associated with an increased risk of complications. *Aust N Z J Med* 1996; 26: 33–39.
43. Cadman A, Lawrance JA, Fitzsimmons L et al. To clot or not to clot? That is the question in central venous catheters. *Clin Radiol* 2004; 59: 349–355.
44. Caers J, Fontaine C, Vinh-Hung V et al. Catheter tip position as a risk factor for thrombosis associated with the use of subcutaneous infusion ports. *Support Care Cancer* 2005; 13: 325–331.
45. Savage KJ, Wells PS, Schulz V et al. Outpatient use of low molecular weight heparin (Dalteparin) for the treatment of deep vein thrombosis of the upper extremity. *Thromb Haemost* 1999; 82: 1008–1010.
46. Rodenhuis S, van't Hek LG, Vlasveld LT et al. Central venous catheter associated thrombosis of major veins: thrombolytic treatment with recombinant tissue plasminogen activator. *Thorax* 1993; 48: 558–559.
47. Schindler J, Bona RD, Chen HH et al. Regional thrombolysis with urokinase for central venous catheter-related thrombosis in patients undergoing high-dose chemotherapy with autologous blood stem cell rescue. *Clin Appl Thromb Hemost* 1999; 5: 25–29.
48. Pucheu A, Dierhas M, Leduc B et al. Fibrinolyse des thromboses veineuses profondes sur dispositifs de perfusion implantables. A propos d'une série consécutive de 57 thromboses et de 32 fibrinolyse. *Bull Cancer* 1996; 83: 293–299.
49. Frank DA, Meuse J, Hirsch D et al. The treatment and outcome of cancer patients with thromboses on central venous catheters. *J Thromb Thrombolysis* 2000; 10: 271–275.
50. Debourdeau P, Elalamy I, de Raignac A et al. Long-term use of daily subcutaneous low-molecular-weight heparin in cancer patients with venous thromboembolism: why hesitate any longer? *Support Care Cancer* 2008; 16: 1333–1341.
51. Buller HR, Agnelli G, Hull RD et al. Antithrombotic therapy for venous thromboembolic disease: the Seventh ACCP Conference on Antithrombotic and Thrombolytic Therapy. *Chest* 2004; 126 (3 Suppl): 401S–428S.